DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: GRANITE	LAKE	Lake Area (ha):	92.19
Town:	STODDARD	Maximum depth (m):	28.9
	Cheshire	Mean depth (m):	9.8
River Basin:		Volume (m³):	9027000
Latitude:		Relative depth:	2.7
Longitude:		Shore configuration:	1.32
Elevation (ft): 1278	Areal water load (m/yr)	: 6.64
Shore length		Flushing rate (yr^{-1}) :	0.70
Watershed are		P retention coeff.:	0.61
% watershed p	onded: 0.0	Lake type: natura	l w/dam

BIOLOGICAL:	10 February 1997	22 July 1996
DOM. PHYTOPLANKTON (% TOTAL) #1	SPARSE - A FEW	SPARSE - NO DOMINANT
#2	DINOBRYON	
#3		
PHYTOPLANKTON ABUNDANCE (units/mL)		
CHLOROPHYLL-A (µg/L)		0.19
DOM. ZOOPLANKTON (% TOTAL) #1	CALANOID COPEPOD 60%	KERATELLA 39%
#2		NAUPLIUS LARVA 23%
#3		CALANOID COPEPOD 18%
ROTIFERS/LITER	<1	28
MICROCRUSTACEA/LITER	5	28
ZOOPLANKTON ABUNDANCE (#/L)	5	56
VASCULAR PLANT ABUNDANCE		Sparse
SECCHI DISK TRANSPARENCY (m)		8.0
BOTTOM DISSOLVED OXYGEN (mg/L)	11.8	8.4
BACTERIA (E. coli, #/100 ml) #1		< 1
#2		< 1
#3		< 1
II		

SUMMER THERMAL STRATIFICATION:

stratified

Depth of thermocline (m): 7.6
Hypolimnion volume (m³): 2260500
Anoxic volume (m³): None

CHEMICAL:			GRANITE I STODDARD	LAKE	
	10 February 1997		22 July 1996		
DEPTH (m)	7.0	14.0	3.5	11.5	21.0
pH (units)	5.6	5.2	6.0	5.4	5.4
A.N.C. (Alkalinity)	1.5	1.0	1.1	0.7	0.9
NITRATE NITROGEN	0.05	0.06	< 0.05		0.09
TOTAL KJELDAHL NITROGEN	< 0.10	< 0.10	< 0.10	< 0.10	0.14
TOTAL PHOSPHORUS	0.005	0.004	0.007	0.006	0.007
CONDUCTIVITY (µmhos/cm)	64.5	64.2	64.6	64.5	66.9
APPARENT COLOR (cpu)	5	5	10	10	10
MAGNESIUM			0.90		
CALCIUM			2.0		
SODIUM			7.0		
POTASSIUM			0.66		
CHLORIDE	10	10	10		11
SULFATE	9	9	8		9
TN : TP					33
CALCITE SATURATION INDEX			5.0		

All results in mg/L unless indicated otherwise


TROPHIC CLASSIFICATION: 1996

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
0	0	0	0	0	Oligo.

COMMENTS:

- 1. Granite Lake was previously surveyed and classified in 1976 and 1985. There was no change in trophic classification or overall water quality over this 20 year period.
- 2. Granite Lake also participates in DES' Volunteer Lake Assessment Program.

 Annual data collected since 1989 show stable or improving water quality.
- 3. Granite Lake is one of only 12 lakes in the state to receive zero trophic points. It continues to maintain its overall superior quality despite the localized impacts from the Route 9 bypass construction in 1994-95.
- 4. Granite Lake is a clear, oligotrophic, moderately acidic lake with very little buffering capacity (A.N.C.). It supports low levels of biological growth (plankton and macrophytes). Chloride and conductivity levels suggest some but well-diluted amounts of salt from highway deicing reaches the lake.



FIELD DATA SHEET

LAKE: GRANITE LAKE TOWN: STODDARD DATE: 07/22/96 WEATHER: CLEAR & SUNNY


DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	21.5	9.0	101 %
1.0	21.5	9.0	101 %
2.0	21.3	9.0	101 %
3.0	21.2	9.0	101 %
4.0	21.0	9.0	101 %
5.0	21.0	9.0	101 %
6.0	21.0	9.0	101 %
7.0	20.0	9.6	104 %
8.0	16.0	12.3	124 %
9.0	14.0	12.5	120 %
10.0	11.4	12.6	113 %
11.0	10.0	12.2	108 %
12.0	8.9	11.6	97 %
13.0	8.0	12.0	101 %
14.0	7.5	11.2	92 %
15.0	6.8	10.6	84 %
17.0	6.3	10.0	80 %
19.0	6.0	9.7	77 %
21.0	5.5	9.5	73 %
23.0	5.5	9.2	72 %
25.0	5.5	8.6	66 %
27.0	5.5	8.4	65 %

SECCHI DISK (m): 8.0 COMMENTS:

BOTTOM DEPTH (m): 27.8

TIME: 1155

*Dissolved oxygen values are in mg/L



AQUATIC PLANT SURVEY

LAK	E: GRANITE LAKE	TOWN: STODDARD	DATE: 07/22/96	
17.0	PLANT	NAME	ABUNDANCE	
Key	GENERIC	COMMON		
М	Myriophyllum humile	Water milfoil	Sparse	
S	Sparganium	Bur reed	Sparse	
P	Pontederia cordata	Pickerelweed	Sparse	
Y	Nuphar	Yellow water lily	Sparse	
G	Gramineae	Grass family	Sparse	
е	Eleocharis	Spike rush	Sparse	
			-	
	·			
		OVERALL ABUNDANCE:	Sparse	

GENERAL OBSERVATIONS:

- 1. Plants continued to be very sparse, although a few more were observed than in 1985.
- 2. The observed milfoil was the native, low-growing variety.
- 3. Previous surveys revealed a 100 foot depth contour. In 1996 the maximum depth sounded was 95 feet. It is assumed the deep spot was missed during the 1996 soundings, but additional soundings should be taken to verify this.